

Inattainability of Carnot efficiency in the Brownian heat engine

Tsuyoshi Hondou*

Department of Physics, Tohoku University
Sendai 980-8578, Japan

Ken Sekimoto†

Yukawa Institute for Theoretical Physics, Kyoto University
Kyoto 606-8502, Japan

We discuss the reversibility of Brownian heat engine. We perform asymptotic analysis of Kramers equation on Büttiker-Landauer system and show quantitatively that Carnot efficiency is unattainable even in a fully overdamping limit. The unattainability is attributed to the inevitable irreversible heat flow over the temperature boundary.

PACS number: 05.40.-a, 05.10.Gg, 05.70.Ln, 87.10.+e

How efficiently can Brownian heat engine work? This question is important not only for the construction of theory of molecular motors [1] but also for foundation of non-equilibrium statistical physics. Like Carnot cycle, Brownian heat engine can extract work from the difference of the temperature of heat baths, where Brownian working material operates as a transducer of thermal energy into mechanical work. The feature of this engine is: 1) It operates autonomously. 2) It is driven by *finite* difference of the temperature of heat baths both of which contact with the working material simultaneously. Thus, this engine works because the system is out of equilibrium. Feynman [2] devised what is called Feynman's ratchet that can rectify the thermal fluctuation for work using the difference of the temperature of two thermal baths. Büttiker [3] and Landauer [4] proposed a simpler type of Brownian motor and pointed out that one could extract work even by the simple heat engine where a Brownian particle is subject to a spatially periodic heat baths in a periodic potential [5].

One crucial point on the Brownian engines is the efficiency [2,6–17]. Feynman claimed that his thermal ratchet can operate reversibly, resulting in Carnot efficiency. Recently, however, some authors claimed that Feynman's claim was incorrect, while some author supported it: Parrondo and Español discussed that Feynman's ratchet should not work reversibly since the engine is simultaneously in contact with heat baths at different temperatures [6]. Sekimoto devised so-called "stochastic energetics" and applied it into Feynman's ratchet [7]. He showed numerically that the efficiency is much less than that of Carnot. Hondou and Takagi showed that reversible operation of Feynman's ratchet is impossible using *reductio ad absurdum* [10]. Magnasco and Stolovitzky studied how the engine generates motion with detailed analysis of its phase space [11]. On the other hand, Sakaguchi claimed that the Feynman's ratchet could operate reversibly by proposing a "stochastic boundary condition" [8]. Similar result is also found in ref. [15] (not

on Feynman's ratchet but on Büttiker-Landauer system), on which detailed discussion will be made later. These studies have reminded us that there is difficulty as to the energetic description on Brownian systems, because naive application of conventional energetics formulated in thermodynamic and/or equilibrium system into Brownian system may lead incorrect result.

Operation of Brownian engines is done by the engines themselves and the engines are, therefore, *out of* equilibrium. To clarify the non-equilibrium nature of Brownian heat engine and to find how we should apply energetics on it, it is important to make a quantitative analysis of the efficiency without adopting over-simplification, for analysis, that loses the function of a heat engine. Because Feynman's ratchet is somewhat complex to make a rigorous analysis, it seems suitable to discuss Büttiker-Landauer [3,4] system that is the simplest system of Brownian motors. Recently, Matsuo and Sasa analyzed the energetics of Büttiker-Landauer system by renormalization method [15]. They claimed the system under quasi-static process approaches Carnot efficiency in overdamping limit [18]. Their analysis was based on a rigorous calculation starting from Kramers equation, and the result is clear except one point: They assumed that the momentum degree of freedom is always in equilibrium with heat bath because the system is overdamping [19]. This assumption is not easy for us to accept because the system is singular at the transition points [20] where the temperature of the heat bath changes suddenly. We conjectured that the essence of the mechanism of Brownian heat engine be concentrated on this singular point and that the nature of these non-equilibrium engines would emerge by the analysis. Thus we will discuss the energetics of Büttiker-Landauer system with paying our attention to the transition points. The result will also give us an insight about how we should apply "stochastic energetics" [7] to the overdamping systems with space-dependent temperature.

Let us consider a one-dimensional Brownian system

*e-mail: hondou@cmpt.phys.tohoku.ac.jp

†e-mail: sekimoto@yukawa.kyoto-u.ac.jp

that Büttiker and Landauer discussed, where working particles operate due to the broken uniformity of the temperature of the heat baths. [3,4]. While Büttiker and Landauer started their discussion from overdamping equation of the system, we start from more basic standpoint of underdamped description, from which the overdamped equation is obtained by eliminating the momentum variable. The probability density in phase space, $\rho(p, q)$, obeys Kramers equation [21]:

$$\begin{aligned} \frac{\partial \rho(p, q)}{\partial t} = & - \left(\frac{\partial J_q}{\partial q} + \frac{\partial J_p}{\partial p} \right) = -K(q) \frac{\partial \rho(p, q)}{\partial p} - \frac{p}{m} \frac{\partial \rho(p, q)}{\partial q} \\ & + \frac{\gamma}{m} \frac{\partial}{\partial p} \left[p \rho(p, q) + m k_B T(q) \frac{\partial \rho(p, q)}{\partial p} \right], \end{aligned} \quad (1)$$

where $K(q) = -\frac{\partial U}{\partial q}$; γ , m , J_q , and J_p are a friction constant, mass of a particle, and probability current in space and that in momentum, respectively [22]. The potential $U(q)$ satisfies, $U(q) = U_r(q) + gq$, where $U_r(q+L) = U_r(q)$, g (> 0) is a gradient of the global slope (load) and L is a period. The temperature has the same spatial period as the potential, $T(q+L) = T(q)$. In this Büttiker-Landauer system, there are two heat baths of which the temperature is T_h (for hot bath) and T_c (for cold bath), respectively. Thus, there are the two transition points in a spatial period where the thermal bath affecting the particle changes. Here we restrict ourselves to the case that $\frac{T_c}{T_h} = O(1)$ for simplicity. The system is known to operate as a molecular engine [3,4,8,15] because the particle can *move* against global gradient of the potential. The globally unidirectional motion was attributed to the difference of the temperatures of the baths, since the hot bath can activate working particle more than the cold bath. Suppose that two working particle *climb* the potential, where one is in a hot bath and the other is in a cold bath. Then the working particle in a hot bath reaches the top of the potential hill more frequently than that in a cold bath, leading to the global motion in the present system. Thus, one can store work in proportion to the probability current. To make energetic analysis, we consider a “replica” particle, of which the energy is $E = p^2/2m + U(q)$. Here, the ensemble average over the replicas corresponds to thermodynamic limit [7].

It was shown that the efficiency of this engine can have Carnot efficiency if the irreversible heat transfer at the transition points is physically negligible. Any Brownian motor is irreversible when they operate with finite probability current. Thus, the operation in Carnot efficiency, if possible, must be in the “stalled state” [15], where the probability current in space disappears, $J_q(q) = 0$ (in an overdamped description) or $\int dp J_q(p, q) = 0$ (in an underdamped description). “Quasi-static” operation requires this “stalled state”. Therefore, we evaluate whether and how the irreversible heat flows at the transition point by solving stationary solution of Kramers equation at the stalled state (Eq.1). For this purpose, we will restrict ourselves to the special region, $q \in [-l_h, l_c]$, around the transition point, $q = 0$, where l_h (l_c) are the

width between the transition point and a proper point on hot (cold) bath that satisfies the following inequality:

$$l_{th} \ll l_x \ll L_x \quad (x = h \text{ or } c), \quad (2)$$

where L_h (L_c) is a width of hot (cold) bath ($L_h + L_c = L$), and l_{th} is a characteristic length scale of the transition region in which the probability density is different from that of thermal equilibrium. The length, l_{th} , is the product of thermal velocity, $v_{th} (\sim \sqrt{k_B T/m})$, and the velocity relaxation time, $\tau (= m/\gamma)$: $l_{th} \sim v_{th} \tau$. The choice of l_x does not alter the following result as long as the inequality, Eq.(2), is satisfied. Although we will discuss only one transition region, the asymptotic behavior does not differ in the other transition region. Hereafter we apply the normalization of the probability density $\rho(p, q)$ as [23]

$$\int_{-\infty}^{\infty} dp \int_{-l_h}^{l_c} dq \rho(p, q) = 1. \quad (3)$$

Now, we will formulate the irreversible heat transfer from a heat bath to the working particle. R.h.s. of Kramers equation (Eq.1) has two parts. The first and the second terms are a Liouville operator on the probability density $\rho(p, q)$ and thus preserve the energy. The last term is what describes the energy transfer between the heat bath and the particle, of which the probability current in momentum space is written: $J_p^{irr} = -\frac{\gamma}{m} \left[p \rho(p, q) + m k_B T(q) \frac{\partial \rho(p, q)}{\partial p} \right]$. Because the probability current disappears, $J_p^{irr} = 0$, for the probability density of equilibrium $\rho(p, q) \propto \exp\{-\frac{p^2/2m + U(q)}{k_B T}\}$, the energy flow through J_p^{irr} can be sufficiently described only where $q \in [-l_h, l_c]$. Average heat transfer from hot bath to the particle per unit time, $\langle \frac{dQ_h}{dt} \rangle$, reads

$$\begin{aligned} \left\langle \frac{dQ_h}{dt} \right\rangle & \sim \int_{-\infty}^{\infty} dp \int_{-l_h}^0 dq \frac{\partial E}{\partial p} J_p^{irr} \\ & = - \int_{-\infty}^{\infty} dp \int_{-l_h}^0 dq \frac{p}{m} \frac{\gamma}{m} \left[p \rho(p, q) + m k_B T(q) \frac{\partial \rho(p, q)}{\partial p} \right]. \end{aligned} \quad (4)$$

By integration by part through momentum space, p , and the property that $\rho(p, q)$ exponentially decreases to zero as $p \rightarrow \pm\infty$, we obtain

$$\begin{aligned} \left\langle \frac{dQ_h}{dt} \right\rangle & = -2 \frac{\gamma}{m} \int_{-\infty}^{\infty} dp \int_{-l_h}^0 dq \left(\frac{p^2}{2m} - \frac{k_B T(q)}{2} \right) \rho(p, q) \\ & \equiv -2 \frac{\gamma}{m} \left\langle \frac{p^2}{2m} - \frac{k_B T_h}{2} \right\rangle_h. \end{aligned} \quad (5)$$

This is the formula of the heat transfer from the hot bath to the particle [24]. When the system is in equilibrium with the heat bath, the heat transfer $\langle \frac{dQ_h}{dt} \rangle$ disappears, because the theory of equipartition requires $\langle p^2/2m \rangle = k_B T/2$. This also shows that the energy exchange between the replica particle and the thermal

bath is dominant only near the thermal transition point, $q = 0$, where the average kinetic energy, $p^2/2m$, deviates from $k_B T/2$.

We remark here how the energy flows around the transition point. As we are analyzing the stalled state, the probability density $\rho(p, q)$ is stationary. Thus, the energy density, $\rho_E(q) = \int_{-\infty}^{\infty} dp (p^2/2m + U(q)) \rho(p, q)$, is stationary. Because there is no work in the stalled state, the conservation of energy requires that $\frac{d\langle Q_h + Q_c \rangle}{dt} = 0$. It shows that the same quantity of the heat from hot bath to the particle flows from the particle to the cold bath: $\langle \frac{dQ_h}{dt} \rangle = -\langle \frac{dQ_c}{dt} \rangle$. It should also be remarked that, in the stalled state, the efficiency vanishes except that the both sides of the last equation vanishes, because the work, the numerator of the efficiency, is absent here. Quasi-static operation is reversible only if the last equation vanishes. Therefore the quantity $\langle \frac{dQ_h}{dt} \rangle$ sufficiently characterizes the operation at the stalled state and thus we will analyze it in detail. Note that the following equation is simultaneously derived [25],

$$\left\langle \frac{dQ_h}{dt} \right\rangle = \int_{-\infty}^{\infty} dp \frac{p^2}{2m} \frac{p}{m} \rho(p, q) \Big|_{q=0}. \quad (6)$$

The formula confirms that the irreversible heat transfer is carried microscopically as a kinetic energy of the particle at a transition point.

It is known, for example by the kinetic theory of gases [26], that there is finite heat transfer, I , in the system where a Brownian particle of finite mass and friction is crossing over the two regions with different temperature, even if the two thermal baths have no direct contact. It implies $I \equiv \langle \frac{dQ_h}{dt} \rangle > 0$. The authors of Ref. [15] assumed that the heat transfer should disappear in overdamping limit, $\frac{m}{\gamma} \rightarrow 0$. However, their assumption is not evident *a priori*. To reveal the validity of the assumption we have to perform proper energetic analysis *on* Kramers equation which includes the degree of momentum, p , instead of overdamped Fokker-Planck equation which lacks the degree.

Hereafter, we will consider the asymptotic behavior of the heat transfer, I , in the limit of overdamping process ($\gamma \rightarrow +\infty$ and/or $m \rightarrow 0$). To find out the asymptotic behavior, it is convenient to use reference heat transfer I^* of unit mass and friction in an arbitrary set of units: $I^* \equiv I(m=1, \gamma=1)$ [27]. By Eq.(5), the reference heat transfer I^* reads:

$$\begin{aligned} I^* &= -2 \left\langle \frac{p^2}{2} - \frac{k_B T_h}{2} \right\rangle_h \\ &= -2 \int_{-\infty}^{\infty} dp \int_{-l_h}^0 dq \left(\frac{p^2}{2} - \frac{k_B T_h}{2} \right) \rho^*(p, q), \end{aligned} \quad (7)$$

where $\rho^*(p, q)$ is a probability density in the reference state: $m = 1$ and $\gamma = 1$. We call a probability density ρ and heat transfer of arbitrary mass and friction in a proper set of units as the generic probability density and the generic heat transfer. Note that the following result is not altered if we have a different reference state. The choice of the values $m = 1$ and $\gamma = 1$ for reference state

is only for simplicity. In the reference state, the characteristic length of the transition region, l_{th}^* , that the probability density in momentum, p , is out of equilibrium is: $l_{th}^* = v_{th}^* \tau^* = \sqrt{k_B T}$.

To evaluate the generic heat transfer (Eq.4) in terms of the reference heat transfer (Eq.4), we will find out the relation between the generic probability density with arbitrary mass and friction $\rho(p, q)$ and the reference one $\rho^*(p, q)$. The potential term $K \frac{\partial \rho}{\partial p}$ of Kramers equation, Eq.(1), is negligible when one discusses the asymptotic behavior of the overdamping [28]. With stationary condition, $\frac{\partial}{\partial t} = 0$, on Eq.(1), we obtain the simple equation that describes stationary flow in phase space around the boundary, $q = 0$:

$$\frac{p}{m} \frac{\partial \rho(p, q)}{\partial q} = \frac{\gamma}{m} \frac{\partial}{\partial p} \left[p \rho(p, q) + m k_B T(q) \frac{\partial \rho(p, q)}{\partial p} \right]. \quad (8)$$

We found here that this equation has a scaling property in mass and friction: The generic probability density $\rho(p, q)$ is expressed using the probability density of the reference state $\rho^*(p, q)$.

$$\rho(p, q) = c \rho^* \left(\frac{p}{\sqrt{m}}, \frac{\gamma}{\sqrt{m}} q \right), \quad (9)$$

where the constant factor c should be determined by normalization (Eq. 3) [29].

As q departs from the transition point, $q = 0$, further than the characteristic length l_{th} , the probability density approaches that of equilibrium, where $\rho_h(p, q) = C_h \exp \left\{ -\frac{p^2}{2m k_B T_h} \right\}$ (for $q \ll -l_{th}$), and $\rho_c(p, q) = C_c \exp \left\{ -\frac{p^2}{2m k_B T_c} \right\}$ (for $q \gg l_{th}$). The coefficients, C_h and C_c , are then required to satisfy the condition of the continuity of probability current. Thus, we have

$$C_h T_h^{3/2} = C_c T_c^{3/2}, \quad (10)$$

which is consistent with the condition derived for the overdamping limit [4]. The remaining condition that determine C_x is the normalization. Note that the normalization of the probability density, ρ , is satisfactorily carried out even by neglecting the contribution from the transition region because the characteristic scale of the transition region l_{th} is much smaller than the width l_x : $l_{th}/l_x \ll 1$ ($x = h$ or c) (Eq.2). Then C_h and C_c are determined as:

$$C_h = \frac{1}{\sqrt{2\pi m k_B T_h}} \frac{T_c}{T_c l_h + T_h l_c}, \quad C_c = \frac{1}{\sqrt{2\pi m k_B T_c}} \frac{T_h}{T_c l_h + T_h l_c}. \quad (11)$$

With these solutions and Eq.(9), we obtain the relation between the two normalized probability densities ρ and ρ^* [29]:

$$\rho(p, q) = \frac{1}{\sqrt{m}} \rho^* \left(\frac{p}{\sqrt{m}}, \frac{\gamma}{\sqrt{m}} q \right). \quad (12)$$

Note that this equation is valid even within the transition region.

We can now express the heat transfer, I , in terms of the reference heat transfer, I^* . We rewrite I as:

$$I = -2\frac{\gamma}{m} \int_{-\infty}^{\infty} dp \int_{-l_h}^0 dq \left(\frac{p^2}{2m} - \frac{k_B T_h}{2} \right) \rho(p, q). \quad (13)$$

By change of variables such that $p' = \frac{p}{\sqrt{m}}$, $q' = \frac{\gamma}{\sqrt{m}}q$ [29], we obtain

$$I = -2 \int_{-\infty}^{\infty} dp \int_{-\frac{\gamma l_h}{\sqrt{m}}}^0 dq \left(\frac{p^2}{2} - \frac{k_B T_h}{2} \right) \rho \left(\sqrt{m}p, \frac{\sqrt{m}q}{\gamma} \right). \quad (14)$$

This yields using Eq.(12),

$$I = -2 \int_{-\infty}^{\infty} dp \int_{-\frac{\gamma l_h}{\sqrt{m}}}^0 dq \left(\frac{p^2}{2} - \frac{k_B T_h}{2} \right) \frac{1}{\sqrt{m}} \rho^*(p, q). \quad (15)$$

This integrand is dominant only near the transition point, $q = 0$, with characteristic length l_{th}^* . As we are analyzing the asymptotic behavior such that $m \rightarrow 0$ and/or $\gamma \rightarrow \infty$, the inequality, $(l_{th}^* \ll) l_h \ll \gamma l_h / \sqrt{m}$, is satisfied. Because the contribution from the interval $q \in [-\gamma l_h / \sqrt{m}, -l_h]$ to the integral is negligible in Eq.(15) compared with that from $q \in [-l_h, 0]$, the interval of this integral may sufficiently be replaced by $q \in [-l_h, 0]$. Using Eq.(7), we obtain one of the main result of our paper [30]:

$$I \sim -\frac{2}{\sqrt{m}} \int_{-\infty}^{\infty} dp \int_{-l_h}^0 dq \left(\frac{p^2}{2} - \frac{k_B T_h}{2} \right) \rho^*(p, q) = \frac{1}{\sqrt{m}} I^*. \quad (16)$$

Because the characteristic length of the transition region vanishes in the overdamping limit, the scaling property is exact asymptotically.

From this result, we can learn that the irreversible heat transfer at the transition point does not decrease when one takes overdamping limit, which is contrast to the claim in Ref. [15]: One way to take this limit is to increase the friction constant, γ : Then, the heat transfer does not decrease, because the heat transfer I does not depend on γ . The other way is to decrease the mass, m : Then, the heat transfer does not decrease neither, moreover the heat transfer increases in the power of $1/\sqrt{m}$. The result justifies the intuitive estimation by Derényi and Astumian [19]. The heat flow is a result of broken symmetry of probability density in momentum at the transition point, because the heat transfer disappears if the probability density is symmetric in phase space, as found by Eq.(6). Since an overdamped equation has no degree of freedom to describe the irreversible flow caused by the discontinuity of the temperature, the previous literature reached Carnot efficiency [15].

Up to now, we have discussed how heat transfer between the two heat baths behaves in the overdamping process. We found that the irreversible heat transfer does not decrease in the process. One finds, however, that the possible work out of the system may also vary according to the overdamping limit, because the probability current may vary due to the change of the parameters, γ and m . Thus, it is not yet obvious whether non-vanishing heat transfer, I , itself reveals that the system cannot attain Carnot efficiency in any condition including non-stalled

state. Thus, in addition to the irreversible heat transfer discussed above, we will estimate work and work-induced heat transfer in an overdamping process.

We will return to the original Kramers equation (Eq.1) for Büttiker-Landauer system. We have analyzed this equation with retaining both degrees of freedom, p and q . However, we do not have to consider a momentum degree of freedom when we discuss work out of the Brownian system, because the work is the function only of the displacement of the position. Thus, we start the evaluation of the work by the overdamped Fokker-Planck equation for the probability density $P(q)$ of the system:

$$\frac{\partial P(q)}{\partial t} = -\frac{\partial}{\partial q} J(q) = \frac{1}{\gamma} \frac{\partial}{\partial q} \left[\frac{\partial U(q)}{\partial q} + \frac{\partial (k_B T(q))}{\partial q} \right] P(q), \quad (17)$$

where the periodic boundary condition is applied: $P(0) = P(L)$ and $\frac{dP}{dq}|_{q=0} = \frac{dP}{dq}|_{q=L}$. Explicit mass dependence on the displacement of the system disappears in the overdamping limit. In stationary state, $\frac{\partial P(q)}{\partial t} = 0$, the probability current, $J(q)$, is independent of q . The probability current J reads

$$J = -\frac{1}{\gamma} \left[\frac{\partial U(q)}{\partial q} + \frac{\partial (k_B T(q))}{\partial q} \right] P(q). \quad (18)$$

The equation for $P(q)$ reads

$$\frac{\partial}{\partial q} \left[\frac{\partial U(q)}{\partial q} + \frac{\partial (k_B T(q))}{\partial q} \right] P(q) = 0. \quad (19)$$

This equation shows that the change of the friction constant, γ , does not alter the probability density, $P(q)$. Thus, with Eq.(18), the probability current, J , scales as: $J \propto \gamma^{-1}$. For a fixed load potential, the work per unit time $\frac{dW}{dt}$ is proportional to its probability current. Thus the sole operation, $\gamma \rightarrow \infty$, does not lead the system to Carnot efficiency, because the induced work ($\propto J$) decreases while the irreversible heat (Eq. (16)) does not decrease.

To find out the mass dependence on the work, we consider the working particle obeying the Stokes' law with its radius r_B , where the mass and the friction are specified by one parameter, r_B : $\gamma \propto r_B$ and $m \propto r_B^3$. Thus, we have $\frac{dW}{dt} \propto J \propto \gamma^{-1} \propto r_B^{-1}$. The irreversible heat transfer $\frac{dQ_{irr}}{dt}$ that is independent of work is just the heat transfer, I (Eq.(16)). Thus we have $\frac{dQ_{irr}}{dt} \propto m^{-1/2} \propto r_B^{-3/2}$. The work-induced heat transfer Q_W that is proportional to the work W is proportional to the probability current J [15]. Thus, we obtain $\frac{dQ_W}{dt} \propto J \propto r_B^{-1}$. The three components determines the efficiency. Thus, we have

$$\eta = \frac{\frac{dW}{dt}}{\frac{dQ_W}{dt} + \frac{dQ_{irr}}{dt}} = \frac{c_1 r_B^{-1}}{c_2 r_B^{-1} + c_3 r_B^{-3/2}} = \frac{c_1}{c_2 + \frac{c_3}{\sqrt{r_B}}}, \quad (20)$$

where c_1 , c_2 and c_3 are constants. The result tells that the efficiency decreases monotonically to zero when one takes overdamping limit $r_B \rightarrow 0$. This result is not altered even if one includes another transition point in the

same period, because the asymptotic behavior of the two are the same.

In this paper, we have analyzed the energetics of a Brownian motor of Büttiker-Landauer type. We showed quantitatively that irreversible heat transfer does not disappear even if one takes overdamping limit ($\gamma \rightarrow +\infty$ and/or $m \rightarrow 0$). This result is in contrast to the claims by Refs. [8,15]. The mass dependence on the irreversible heat is consistent to the intuitive estimation by Ref. [19]. We further analyzed the effect of non-vanishing irreversible heat transfer on the efficiency and showed that, even in fully overdamping limit, Carnot efficiency is inattainable for the particle obeying Stokes' law. It shows that the maximum efficiency of the Brownian motor is not attained in the stalled state. The result revealed that the Brownian heat engine is qualitatively different from heat engines of which the most efficient operation is in quasi-static: Quasi-static process is the worst condition for the Brownian heat engine to work, while is the best for Carnot cycle.

The location of irreversible heat transfer is the transition region characterized by the thermal length l_{th} . It is certain that the characteristic length l_{th} may disappear in the fully overdamping limit. Then, however, the irreversible effect at the transition region cannot be eliminated. From the result we also learn a lesson how to apply energetics to overdamping systems with a space-dependent temperature: We should apply energetics *before* taking overdamping limit. Otherwise, we might fail in proper evaluation of irreversible heat transfer within the transition region [8,15], because energetic interaction between the heat bath and the particle is carried by the momentum exchange between them. When the particle has smaller kinetic energy than that expected by equipartition theorem, the particle receive kinetic energy from the heat bath in average. Thus, if we lose the degree of momentum as in the overdamped equation, we cannot describe this existing physical process properly.

The present system cannot have maximum efficiency at a quasistatic condition. This means that the maximum efficiency is achieved with finite probability current, which is therefore accompanied by irreversible dissipation. Thus, the next challenging question would be, "Is there any principle that determines the optimal efficiency in Brownian heat engines?"

We would like to thank T. Chawanya, T. Mizuguchi, H. Hayakawa, H. Nishimori, M. Matsuo, S. Sasa, T. Shibata, Y. Hayakawa, M. Sano, F. Takagi, Y. Sawada, and T. Tsuzuki for helpful discussion. This work is supported in part by the Japanese Grant-in-Aid for Science Research Fund from the Ministry of Education, Science and Culture (Nos. 09740301, 11156216 and 12740226) and the Inamori Foundation.

[1] See for review: F. Jülicher, A. Ajdari and J. Prost, Rev. Mod. Phys. **69**, 1269 (1997).

- [2] R. P. Feynman, R. B. Leighton and M. Sands: *The Feynman Lectures in Physics* (Addison-Wesley, Massachusetts, 1966).
- [3] M. Büttiker, Z. Phys. B **68**, 161 (1987).
- [4] R. Landauer, J. Stat. Phys. **53**, 233 (1998).
- [5] Energetics was not discussed quantitatively at that time, because little was known in the energetic evaluation of Brownian systems.
- [6] J. M. R. Parrondo and P. Español, Am. J. Phys. **64**, 1125 (1996).
- [7] K. Sekimoto: J. Phys. Soc. Jpn. **66**, 1234 (1997).
- [8] H. Sakaguchi, J. Phys. Soc. Jpn. **67**, 709 (1998).
- [9] H. Kamegawa, T. Hondou, and F. Takagi, Phys. Rev. Lett. **80**, 5251 (1998).
- [10] T. Hondou and F. Takagi, J. Phys. Soc. Jpn. **67**, 2974 (1998).
- [11] M. O. Magnasco and G. Stolovitzky, J. Stat. Phys. **93**, 615 (1998).
- [12] T. Shibata and S. Sasa, J. Phys. Soc. Jpn. **67**, 2666 (1998).
- [13] J.M.R. Parrondo, Europhys. Lett. **43**, 248 (1998).
- [14] K. Sekimoto, J. Phys. Soc. Jpn. **68**, 1448 (1999).
- [15] M. Matsuo and S. Sasa, Physica A **276**, 188 (2000).
- [16] A. Parmeggiani *et al.*, Phys. Rev. E **60**, 2127 (1999).
- [17] F. Takagi and T. Hondou, Phys. Rev. E **60**, 4954 (1999).
- [18] The similar claim and assumption is found in Ref. [8].
- [19] In the coarse of our preparation of manuscript, we found the article: I. Derényi and R.D. Astumian, Phys. Rev. E **59**, 6219 (1999).
- [20] The term "transition" was introduced in Ref. [4] in the sense that the probability density in overdamping limit changes suddenly.
- [21] H.A. Kramers, Physica **7**, 284 (1940).
- [22] In this paper, we discussed the case that friction coefficient γ is independent of the temperature T for simplicity. Friction coefficient may depend on the temperature as discussed in Ref. [4]. However, it does not alter the central result, because asymptotic property in overdamping process is the same for temperature dependent friction coefficient.
- [23] The choice of the interval, $q \in [-l_h, l_c]$, changes the factor of the probability density $\rho(p, q)$. However, the choice does not alter the following asymptotic estimation in an overdamping process.
- [24] This formula is consistent with that derived in a different way in Ref. [6].
- [25] The equation can be obtained even just *at* the transition point, because $\rho(p, q)$ is continuous.
- [26] See, for example, G.M. Barrow, Physical Chemistry 4th ed. (McGraw-Hill, New York 1979).
- [27] We require the reference state to hold Eq.(2): $l_{th}^* \ll l_h \ll L_h$.
- [28] If one takes overdamping limit, characteristic length of the transition region, l_{th} , decreases to zero. Thus, the effect of potential force, K , in the transition region disappears in comparison to the interaction between heat bath and the system.
- [29] Here, the parameters, m and γ , are the ratios of those between in the generic state and in the reference state. Thus, they are dimensionless.
- [30] The same result can also be obtained using Eq.(6) instead of Eq.(5).